

**Marc-Gilbert Lagny
Francine Blaffart
ECCP
Department of Nursing and
Department of Cardiovascular and Thoracic Surgery
Prof. JO Defraigne
University Hospital of Liège, Belgium.**




CPB AND AORTIC SURGERY

The State of the Art

From a Theoretical to a Practical Approach



M-G LAGNY, F. BLAFFART, ECCP CHU de Liège



INTRODUCTION

- Prevention

Michel JB, et al.

Novel aspects of the pathogenesis of aneurysms on the abdominal aorta in humans. Cardiovasc Res. 2011 Apr 1;90(1):18-27

Golledge J, Norman PE,

- Medical treatment

Current status of medical management for abdominal aortic aneurysm. Atherosclerosis, 2011. 217 (1):p.57-63.

- Endovascular aortic repair

Nienaber CA, et al.

Randomized comparison of strategies for type B aortic dissection: the INvestigation of STEnt Grafts in Aortic Dissection (INSTEAD) trial. Circulation, 2009. 120(25):p.2519-28.


Hao Z, et al.

Endovascular stent-graft placement or open surgery for the treatment of acute type B aortic dissection: a meta-analysis. Ann Vasc Surg, 2012. 26(4):p.454-61.

AIM OF THE TOPIC




CPB for aortic surgery: state of the art



- Knowledge
- Anatomical and pathophysiological pre-requests
- Flexibility
- Equipment

Ascending aortic surgery and arterial cannulation:



**Surgical
repairment**

Ascending aorta

**Left subclavian/
axillary**

Femoral

Other

The arterial cannulation



Cannulation site	Advantages	Disadvantages
Femoral artery	Ease of access Size Adequate flow rates	Retrograde flow Malperfusion Proximal embolization
Axillary/subclavian artery	Reports of reduced mortality and stroke	Arm numbness Brachial plexus injury
Aortic cannulation	Speed of cannulation	Embolic event
Ventricular apex	Adequacy of flow	Ventricular injury
Direct	Antegrade flow Direct cannulation of true lumen	

eTable 3: The potential advantages and disadvantages of the different cannulation sites utilized in acute type A aortic dissection

Bonser RS, et al.
Acute Aortic Dissection. JAAC Vol. 58, No. 24, 2011.

Arterial cannulation complication and perfusion: local dissection

- Pressure monitoring




SWITCH TO ANOTHER CANNULATION SITE

Anticipation: Y Line

Arterial cannulation complication and perfusion: malperfusion (FLAP)

- Sudden increase of arterial line pressure
- Inequat cerebral perfusion?: NIRS, TCD
- Inadequate spinal cord perfusion (MEP)
- Late diagnostic (lactates)
- Left radial pressure




**Pre and post arch arterial lines
(+ left femoral)
Switch to another cannulation site
RE-INSTORE ANTEGRADE FLOW**



Arterial femoral cannulation complication and perfusion: false lumen


- Pressure monitoring
- Transesophageal echocardiography (TOE)
- Arterial pressure (left radial)



CHECK THE CANNULATION
Switch to another cannulation site

Arterial cannulation complication and perfusion: embolic event

- Doppler
- Specific cannula



Christenson JT, et al.
Tex Heart Inst J. 2005;32(4):515-21.



Cerebral protection

Protecting the brain during aortic surgery: an enduring debate with unanswered questions.


Stein LH, Elefteriades JA,

Section of Cardiothoracic Surgery, Yale University School of Medicine, New Haven, CT 06510, USA.


Cardiothorac Vasc Anesth. 2010 Apr;24(2):316-21. Epub 2009 Jul 30.




Surgery on the ascending aorta and the arch cerebral protection





Selective cerebral perfusion:



Antegrade selective cerebral perfusion: Complications and monitoring

Pro	Cons	Monitoring
Control of flow delivery	Local dissection Downstream dissection	Pression Nirs, Doppler
	Embolic load	Doppler
	Cerebral oedema in case of overflow and or over pressure	Flow and pressure control
	Integrity of the circle of Willis? in case of single carotid perfusion	NIRS, transcranial doppler, Left radial arterial pressure (60mmHg) (JbSVO ₂) (S100 protein; NSE)

Retrograde cerebral perfusion: complications and monitoring


Pro	Cons	monitoring
Easy of access	Poor control of flow delivery, Dispersion of the flow through the Azygos vein	NIRS, transcranial doppler
Retrograde flush of the carotids	Cerebral oedema in case of overflow And or overpressure	Flow control and Venous pressure (30 mmHg)
<div>   </div>		

Deep hypothermia circulatory arrest: state of the art



- Respect of temperatures gradients (6-10°C max)
- Normoxia
- Hct level versus viscosity (25% Hct max)
- Homogenization of temperatures (cerebral and systemic)
- Hardware:
 - Heater cooler device
 - Efficient heat exchanger
 - Cooling helmet
 - Blanket


Deep hypothermia circulatory arrest: state of the art: blood gases management



Deep hypothermia circulatory arrest euglycemia

Avoiding Stroke During Cardiac Surgery

Kristine Kellermann, DVM¹, and Bettina Jungwirth, MD¹

Seminars in Cardiothoracic and
Vascular Anesthesia
14(2) 95–101
© The Author(s) 2010
Reprints and permission: <http://www.sagepub.com/journalsPermissions.nav>
DOI: 10.1177/1089253210370902
<http://scv.sagepub.com>


Abstract

The life saving benefits of cardiac surgery are frequently accompanied by negative side effects such as stroke, that occurs with an incidence of 2%-13% dependent to type of surgery. The etiology is most likely multifactorial with embolic events considered as main contributor. Although stroke presents a common complication, no guidelines for any routine use of pharmacological substances or non-pharmacological strategies exist to date.

Non-pharmacological strategies include monitoring of brain oxygenation and perfusion with devices such as near infrared spectroscopy and Transcranial Doppler help. Epiaortic and transesophageal echocardiography visualize aorta pathology, enabling the surgeon to sidestep atheromatous segments. Additionally can the use of specially designed aortic cannulae and filters help to reduce embolization. Brain perfusion can be improved by using antero- or retrograde cerebral perfusion during deep hypothermic circulatory arrest, by tightly monitoring mean arterial blood pressure and hemodilution. Controlling perioperative temperature and glucose levels may additionally help to ameliorate secondary damage.

Many pharmacological compounds have been shown to be neuroprotective in preclinical models, but clinical studies failed to confirm these results so far.

Remacemide, an NMDA-receptor-antagonist showed a significant drug-based neuroprotection during cardiac surgery. Other substances currently assessed in clinical trials whose results are still pending are acadesine, an adenosine-regulating substance, the free radical scavenger edaravone and the local anesthetic lidocaine.

Stroke remains as significant complication after cardiac surgery. Non-pharmacological strategies allow perioperative caregivers to detect injurious events and to ameliorate stroke and its sequelae. Considering the multi-factorial etiology though, stroke prevention will likely have to be addressed with an individualistic combination of different strategies and substances.

Deep hypothermia circulatory arrest and reperfusion injury

- Low pressure
- Normoxia
- Reperfusion solution?


Deep hypothermic circulatory arrest and global reperfusion injury: Avoidance by making a pump prime perfusate—A new concept

Bradley S. Allen, MD

J Thorac Cardiovasc Surg 2003; 125:625-32

- Hyperkalemia (?) → hemodiafiltration

Descending aortic surgery



**Surgical
repair**

**Left-left
Bypass**

**Right-left
Bypass**

**Conventional
CPB**



None

Descending aortic surgery: CPB circuit

Left-left bypass Left-right bypass	Conventional miniaturized CPB
Left atrium – distal aorta Right atrium distal aorta	Right atrium (femoral access) – distal aorta
Centrifugal pump Autoregulation of the volemia	Centrifugal or roller pump
	Heat exchanger + oxygenator
Low heparin level	Full heparinized
	Easy shunt for selective perfusion
	Quick response to acute hemorrhagic event

Surgery on the descending aorta medullar and splanchnic selective perfusion

- Perfusion
- Local hypothermia
- Systemic mild hypothermia (32°C)



Medullar and splanchnic perfusion: complications and monitoring

Complications	monitoring
Local dissection	Q-Pressure
oedema in case of overflow and or over pressure or brain herniation	CSF drainage (10mmHg)
Malperfusion Upstream embolism	Flow, regional pressure (60mmHg), Doppler flowmetry MEP (motor evoked potential). Mucosal pH tonometry NIRS ?


Coagulation management

Thromboelastometry-guided administration of fibrinogen concentrate for the treatment of excessive intraoperative bleeding in thoracoabdominal aortic aneurysm surgery

Niels Rahe-Meyer, MD, MSc, PhD,^a Cristina Solomon, MD,^a Michael Winterhalter, MD,^a Siegfried Piepenbrock, MD,^a Kenichi Tanaka, MD, MSc, PhD,^b Axel Haverich, MD,^c and Maximilian Pichlmaier, MD^c

Blood management

- Selective suction blood management
- Specific filtration
- Cell saving process
- Selective allogenic blood component transfusion



Conclusions

- Aorta surgery is a team work
- Multiple modal approaches
- Engering developments
- EBM and EBP



**Thank you for your
attention**